



Partial Averaging of Fuzzy Hyperbolic Differential Inclusions

Tatyana Alexandrovna Komleva¹, Irina Vladimirovna Molchanyuk²,
Andrej Viktorovich Plotnikov^{2,*}, Liliya Ivanovna Plotnikova³

¹Department of Mathematics, Odessa State Academy of Civil Engineering and Architecture, Odessa, Ukraine

²Department of Applied Mathematics, Odessa State Academy of Civil Engineering and Architecture, Odessa, Ukraine

³Department of Mathematics, Odessa National Polytechnic University, Odessa, Ukraine

Email address:

a-plotnikov@ukr.net (A.V. Plotnikov)

*Corresponding author

To cite this article:

Tatyana Alexandrovna Komleva, Irina Vladimirovna Molchanyuk, Andrej Viktorovich Plotnikov, Liliya Ivanovna Plotnikova. Partial Averaging of Fuzzy Hyperbolic Differential Inclusions. *International Journal of Systems Science and Applied Mathematics*. Vol. 1, No. 4, 2016, pp. 42-49. doi: 10.11648/j.ijssam.20160104.12

Received: September 19, 2016; **Accepted:** September 28, 2016; **Published:** October 19, 2016

Abstract: In this article, we considered the fuzzy hyperbolic differential inclusions (fuzzy Darboux problem), introduced the concept of R-solution and proved the existence of such a solution. Also the substantiation of a possibility of application of partial averaging method for hyperbolic differential inclusions with the fuzzy right-hand side with the small parameters is considered.

Keywords: Hyperbolic Differential Inclusion, Fuzzy, Averaging, R-solution

1. Introduction

In 1990 J.P. Aubin [6] and V.A. Baidosov [7, 8] introduced differential inclusions with the fuzzy right-hand side. Their approach is based on usual differential inclusions. E. Hüllermeier [20, 21] introduced the concept of R-solution similarly how it has been done in [34]. Later, the various properties of fuzzy solutions of differential inclusions, and their use in modeling various natural science processes were considered (see [1, 4, 5, 17, 18, 26, 27] and the references therein).

The averaging methods combined with the asymptotic representations (in Poincaré sense) began to be applied as the basic constructive tool for solving the complicated problems of analytical dynamics described by the differential equations. After the systematic researches done by N. M. Krylov, N. N. Bogoliubov, Yu. A. Mitropolsky etc, in 1930s, the averaging method gradually became one of the classical methods in analyzing nonlinear oscillations (see [10, 25, 40, 42] and references therein). In works [36-39], the possibility of application of schemes of full and partial averaging for fuzzy differential inclusions with a small parameter was proved.

In papers [2, 3, 9, 11, 13, 18, 22, 30, 32, 33, 35], authors investigate classical models of partial differential equations with uncertain parameters, considering the parameters as

fuzzy numbers. It was an obvious step in the mathematical modeling of physical processes. Study of fuzzy partial differential equations means the generalization of partial differential equations in fuzzy sense. While doing modelling of real situation in terms of partial differential equation, we see that the variables and parameters involve in the equations are uncertain (in the sense that they are not completely known or inexact or imprecise). Many times common initial or boundary condition of ambient temperature is a fuzzy condition since ambient temperature is prone to variation in a range. We express this impreciseness and uncertainties in terms of fuzzy numbers. So we come across with fuzzy partial differential equations. Also obviously, these equations can be written in as fuzzy partial differential inclusions.

In this work we consider fuzzy hyperbolic differential inclusions (fuzzy Darboux problem) and introduce the concept of R-solution similarly how it has been done in [36, 40, 50, 52, 53]. Also we ground the possibility of application of partial averaging method for fuzzy Darboux problem. This result generalizes the results of A. N. Vityuk [40, 52] for the ordinary hyperbolic differential inclusions and M. Kiselevich [23], D. G. Korenevskii [24] for the ordinary hyperbolic differential equations.

2. Preliminaries

Let $\text{comp}(R^n)(\text{conv}(R^n))$ be a family of all nonempty (convex) compact subsets from the space R^n with the Hausdorff metric

$$h(A, B) = \min_{r \geq 0} \{B \subset S_r(A), A \subset S_r(B)\},$$

where $A, B \in \text{comp}(R^n)$, $S_r(A)$ is r -neighborhood of set A .

Let E^n be a family of all $u: R^n \rightarrow [0, 1]$ such that u satisfies the following conditions:

- 1) u is normal, i.e. there exists an $x_0 \in R^n$ such that $u(x_0) = 1$;
- 2) u is fuzzy convex, i.e. $u(\lambda x + (1-\lambda)y) \geq \min\{u(x), u(y)\}$ for any $x, y \in R^n$ and $0 \leq \lambda \leq 1$;
- 3) u is upper semicontinuous, i.e. for any $x_0 \in R^n$ and $\varepsilon > 0$ where exists $\delta(x_0, \varepsilon) > 0$ such that $u(x) < u(x_0) + \varepsilon$ whenever $\|x - x_0\| < \delta(x_0, \varepsilon)$, $x \in R^n$;
- 4) the closure of the set $\{x \in R^n : u(x) > 0\}$ is compact.

If $u \in E^n$, then u is called a fuzzy number, and E^n is said to be a fuzzy number space.

Definition 1. The set $\{x \in R^n : u(x) \geq \alpha\}$ is called the α -level $[u]^\alpha$ of a fuzzy number $u \in E^n$ for $0 < \alpha \leq 1$. The closure of the set $\{x \in R^n : u(x) > 0\}$ is called the 0-level $[u]^0$ of a fuzzy number $u \in E^n$.

It is clearly that the set $[u]^\alpha \in \text{conv}(R^n)$ for all $0 \leq \alpha \leq 1$.

Theorem 1. (Stacking Theorem [31]) If $u \in E^n$ then

- 1) $[u]^\alpha \in \text{conv}(R^n)$ for all $\alpha \in [0, 1]$;
- 2) $[u]^{\alpha_2} \subset [u]^{\alpha_1}$ for all $0 \leq \alpha_1 \leq \alpha_2 \leq 1$;
- 3) if $\{\alpha_k\}$ is a nondecreasing sequence converging to $\alpha > 0$, then $[u]^\alpha = \bigcap_{k \geq 1} [u]^{\alpha_k}$.

Conversely, if $\{A_\alpha : \alpha \in [0, 1]\}$ is the family of subsets of R^n satisfying conditions 1) - 3) then there exists $u \in E^n$ such that $[u]^\alpha = A_\alpha$ for $0 < \alpha \leq 1$ and $[u]^0 = \bigcup_{0 < \alpha \leq 1} A_\alpha \subset A_0$.

Let θ be the fuzzy number defined by $\theta(x) = 0$ if $x \neq 0$ and $\theta(0) = 1$.

Define $D: E^n \times E^n \rightarrow [0, \infty)$ by the relation

$$D(u, v) = \sup_{0 \leq \alpha \leq 1} h([u]^\alpha, [v]^\alpha).$$

Then D is a metric in E^n . Further we know that [41]:

- i). (E^n, D) is a complete metric space,
- ii). $D(u + w, v + w) = D(u, v)$ for all $u, v, w \in E^n$,
- iii). $D(\lambda u, \lambda v) = |\lambda| D(u, v)$ for all $u, v \in E^n$ and $\lambda \in R$.

3. Fuzzy Hyperbolic Differential Inclusion. R-solution

Consider the fuzzy hyperbolic differential inclusion (or in other words, fuzzy Darboux problem)

$$\begin{aligned} u_{xy}(x, y) &\in F(x, y, u(x, y)), \\ u(x, 0) &= \varphi(x), x \in [0, a], \\ u(0, y) &= \psi(y), y \in [0, b], \varphi(0) = \psi(0), \end{aligned} \quad (1)$$

where $u \in R^n, u_{xy}(x, y) = \frac{\partial^2 u(x, y)}{\partial x \partial y}, F: [0, a] \times [0, b] \times R^n \rightarrow E^n, \varphi: [0, a] \rightarrow R^n, \psi: [0, b] \rightarrow R^n$.

We interpret fuzzy Darboux problem (1) as a family of set-valued Darboux problems

$$\begin{aligned} u_{xy}^\alpha(x, y) &\in [F(x, y, u^\alpha(x, y))]^\alpha, \\ u^\alpha(x, 0) &= \varphi(x), x \in [0, a], \\ u^\alpha(0, y) &= \psi(y), y \in [0, b], \\ \varphi(0) &= \psi(0), \alpha \in [0, 1]. \end{aligned} \quad (2)$$

Qualitative properties and structure of the set of solutions of the set-valued Darboux problem have been studied by many authors, for instance [12, 14-16, 28, 29, 40, 44-53] and references therein.

Definition 2 [28, 43]. A function $u: [0, a] \times [0, b] \rightarrow R^n$ is said to be absolutely continuous on $[0, a] \times [0, b]$ ($u(\cdot, \cdot) \in AC([0, a] \times [0, b])$) if there exist absolutely continuous functions $\varphi: [0, a] \rightarrow R^n$ and $\psi: [0, b] \rightarrow R^n$, and Lebesgue integrable function $g: [0, a] \times [0, b] \rightarrow R^n$ such that

$$u(x, y) = \varphi(x) + \psi(y) - c + \int_0^x \int_0^y g(\xi, \zeta) d\zeta d\xi, \varphi(0) = \psi(0) = c.$$

Definition 3. An α -solution $u^\alpha(\cdot, \cdot)$ of (1) is understood to be an absolutely continuous function $u^\alpha: [0, a] \times [0, b] \rightarrow R^n$ that satisfies (2) for almost every $(x, y) \in [0, a] \times [0, b]$ and the boundary conditions for any $x \in [0, a]$ and $y \in [0, b]$.

Let U^α denote the α -solution set of (2) and $U^\alpha(x, y) = \{u^\alpha(x, y) : u^\alpha(\cdot, \cdot) \in U^\alpha\}$. Clearly a family of subsets $U(x, y) = \{U^\alpha(x, y) : \alpha \in [0, 1]\}$ may not satisfy to conditions of Theorem 1, i.e. $U(x, y) \notin E^n$. For example,

$U^\alpha(x, y) \in \text{comp}(R^n)$ and $U^\alpha(x, y) \notin \text{conv}(R^n)$ for any $\alpha \in [0, 1]$. Therefore, we introduce the definition of R-solutions for fuzzy Darboux problem (1).

$$\sup_{\alpha \in [0, 1]} h([R(t + \sigma, s + \eta)]^\alpha, \bigcup_{u \in [R(t, s)]^\alpha} \left\{ v(t + \sigma) + \tau(s + \eta) - u + \int_t^{t+\sigma} \int_s^{s+\eta} [F(\xi, v, u)]^\alpha dv d\xi \right\} \Bigg) = o(\sigma\eta), \quad (3)$$

is called the R-solution of fuzzy Darboux problem (1), where $v(x) \in AC(t, t + \sigma)$, $\tau(y) \in AC(s, s + \eta)$, $v(x) \in [R(x, s)]^\alpha$, $x \in [t, t + \sigma]$, $\tau(y) \in [R(t, y)]^\alpha$, $y \in [s, s + \eta]$,

$$v(t) = \tau(s) = u, \quad \lim_{\substack{\sigma \rightarrow 0_+ \\ \eta \rightarrow 0_+}} \frac{o(\sigma\eta)}{\sigma\eta} = 0.$$

Now we are interested in the following question: Under what conditions, there exists a unique R-solution to (1). In the next theorem we find the existence result for a unique R-solution of fuzzy Darboux problem (1).

Theorem 2. Suppose the following conditions hold:

- 1) fuzzy mapping $F(\cdot, \cdot, u)$ is measurable, for all $u \in R^n$;
- 2) there exists $\lambda > 0$ such that for all $u', u'' \in R^n$

$$D(F(x, y, u'), F(x, y, u'')) \leq \lambda \|u' - u''\|$$

for every $(x, y) \in [0, a] \times [0, b]$;

- 3) there exists $\gamma > 0$ such that $D(F(x, y, u), \theta) \leq \gamma$ for

$$u^\alpha(x_2, y_2) - u^\alpha(x_2, y_1) - u^\alpha(x_1, y_2) + u^\alpha(x_1, y_1) \in \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, u^\alpha(x, y))]^\alpha dy dx$$

for every $[x_1, x_2] \times [y_1, y_2] \subset [0, a] \times [0, b]$.

Consider any solutions $u_1^\alpha(\cdot, \cdot), u_2^\alpha(\cdot, \cdot) \in U^\alpha$ and any $\beta \in [0, 1]$. Let $u_\beta^\alpha(\cdot, \cdot)$ be such that

$$u_\beta^\alpha(x, y) = \beta u_1^\alpha(x, y) + (1 - \beta) u_2^\alpha(x, y)$$

for every $(x, y) \in [0, a] \times [0, b]$.

Then

$$\begin{aligned} & u_\beta^\alpha(x_2, y_2) - u_\beta^\alpha(x_2, y_1) - u_\beta^\alpha(x_1, y_2) + u_\beta^\alpha(x_1, y_1) = \\ & = \beta u_1^\alpha(x_2, y_2) - \beta u_1^\alpha(x_2, y_1) - \beta u_1^\alpha(x_1, y_2) + \beta u_1^\alpha(x_1, y_1) + \\ & + (1 - \beta) u_2^\alpha(x_2, y_2) - (1 - \beta) u_2^\alpha(x_2, y_1) - (1 - \beta) u_2^\alpha(x_1, y_2) + (1 - \beta) u_2^\alpha(x_1, y_1) \in \\ & \in \beta \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, u_1^\alpha(x, y))]^\alpha dy dx + (1 - \beta) \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, u_2^\alpha(x, y))]^\alpha dy dx \subset \\ & \subset \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, \beta u_1^\alpha(x, y) + (1 - \beta) u_2^\alpha(x, y))]^\alpha dy dx = \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, u_\beta^\alpha(x, y))]^\alpha dy dx \end{aligned}$$

Definition 3. The upper semicontinuous fuzzy mapping $R : [0, a] \times [0, b] \rightarrow E^n$ that satisfies to the following system

every $(x, y, u) \in [0, a] \times [0, b] \times R^n$;

- 4) for all $\beta \in [0, 1], u', u'' \in R^n$ and every $(x, y) \in [0, a] \times [0, b]$

$$\beta F(x, y, u') + (1 - \beta) F(x, y, u'') \subset F(x, y, \beta u' + (1 - \beta) u'');$$

- 5) functions $\varphi(\cdot)$ and $\psi(\cdot)$ are absolutely continuous functions on $[0, a]$ and $[0, b]$.

Then there exists a unique R-solution $R(\cdot, \cdot)$ of fuzzy Darboux problem (1) defined on the set $[0, a] \times [0, b]$.

Proof. By [29, 49], every set-valued Darboux problem of family (2) has solution $u^\alpha(\cdot, \cdot)$ on the set $[0, a] \times [0, b]$, i.e. $U^\alpha(x, y) \neq \emptyset$ for every $(x, y) \in [0, a] \times [0, b]$ and $\alpha \in [0, 1]$.

Also by [40] and [50], $U^\alpha(x, y) \in \text{comp}(R^n)$ for every $(x, y) \in [0, a] \times [0, b]$ and $\alpha \in [0, 1]$.

By [40] and [51], if $u^\alpha(\cdot, \cdot) \in U^\alpha$ then

i. e. $u_\beta^\alpha(x_2, y_2) - u_\beta^\alpha(x_2, y_1) - u_\beta^\alpha(x_1, y_2) + u_\beta^\alpha(x_1, y_1) \in \int_{x_1}^{x_2} \int_{y_1}^{y_2} [F(x, y, u_\beta^\alpha(x, y))]^\alpha dy dx$ for every $(x, y) \in [0, a] \times [0, b]$ and $\beta \in [0, 1]$.

By [40] and [51], function $u_\beta^\alpha(\cdot, \cdot)$ is solution of set-valued Darboux problem (2), i.e. $u_\beta^\alpha(x, y) \in U^\alpha(x, y)$ for every $(x, y) \in [0, a] \times [0, b]$. Consequently $U^\alpha(x, y) \in \text{conv}(R^n)$ for every $(x, y) \in [0, a] \times [0, b]$ and $\alpha \in [0, 1]$.

Since, $[F(x, y, u)]^{\alpha_2} \subset [F(x, y, u)]^{\alpha_1}$ for all $0 \leq \alpha_1 < \alpha_2 \leq 1$ and $(x, y, u) \in [0, a] \times [0, b] \times R^n$, then $U^{\alpha_2}(x, y) \subset U^{\alpha_1}(x, y)$ for all $0 \leq \alpha_1 < \alpha_2 \leq 1$ and $(x, y) \in [0, a] \times [0, b]$.

By [50, 52], every Darboux problem of family (2) has one R-solution $R^\alpha(\cdot, \cdot)$ on the set $[0, a] \times [0, b]$ and we have $R^\alpha(x, y) = U^\alpha(x, y)$ for every $\alpha \in [0, 1]$ and $(x, y) \in [0, a] \times [0, b]$.

By [20, 53], we get that a family of subsets $R(x, y) = \{R^\alpha(x, y) : \alpha \in [0, 1]\}$ satisfies to conditions of Theorem 1, i.e. $R(x, y) \in E^n$ for every $(x, y) \in [0, a] \times [0, b]$. This concludes the proof.

4. The Method of Partial Averaging

Now consider fuzzy Darboux problem with the small parameters

$$\begin{aligned} u_{xy}(x, y) &\in \varepsilon_1 \varepsilon_2 F(x, y, u(x, y)), \\ u(x, 0) &= \varphi(x), x \in R_+, \\ u(0, y) &= \psi(y), y \in R_+, \varphi(0) = \psi(0), \end{aligned} \quad (4)$$

where $\varepsilon_1 > 0, \varepsilon_2 > 0$ - small parameters, $R_+ = [0, +\infty)$.

In this work, we associate with the problem (4) the following full averaged fuzzy Darboux problem

$$\begin{aligned} z_{xy}(x, y) &\in \varepsilon_1 \varepsilon_2 G(x, y, z(x, y)), \\ z(x, 0) &= \varphi(x), x \in R_+, \\ z(0, y) &= \psi(y), y \in R_+, \varphi(0) = \psi(0), \end{aligned} \quad (5)$$

where $G : R^n \rightarrow E^n$ such that

$$\lim_{\substack{T_1 \rightarrow \infty \\ T_2 \rightarrow \infty}} D \left(\frac{1}{T_1 T_2} \int_0^{T_1} \int_0^{T_2} F(x, y, z) dy dx, \frac{1}{T_1 T_2} \int_0^{T_1} \int_0^{T_2} G(x, y, z) dy dx \right) = 0. \quad (6)$$

The main theorem of this section is on averaging for fuzzy Darboux problem with the small parameters. It establishes nearness of R-solutions of (4) and (5), and reads as follows.

Theorem 3. Let in the domain $Q = \{(x, y, u) : x \in R_+, y \in R_+, u \in B \subset R^n\}$ the following conditions hold:

- 1) fuzzy mappings $F(\cdot, \cdot, u)$ and $G(\cdot, \cdot, u)$ is continuous on $R_+ \times R_+$;
- 2) fuzzy mappings $F(x, y, \cdot)$ and $G(x, y, \cdot)$ satisfy a

Lipschitz condition

$$\begin{aligned} D(F(x, y, u'), F(x, y, u'')) &\leq \lambda \|u' - u''\|, \\ D(G(x, y, u'), G(x, y, u'')) &\leq \lambda \|u' - u''\| \end{aligned}$$

with a Lipschitz constant $\lambda > 0$;

3) there exists $\gamma > 0$ such that

$$D(F(x, y, u), \theta) \leq \gamma, \quad D(G(x, y, u), \theta) \leq \gamma$$

for every $(x, y) \in R_+ \times R_+$ and every $u \in R^n$;

4) for all $\beta \in [0, 1], u', u'' \in R^n$ and every $(x, y) \in R_+ \times R_+$

$$\beta F(x, y, u') + (1 - \beta) F(x, y, u'') \subset F(x, y, \beta u' + (1 - \beta) u''),$$

$$\beta G(x, y, u') + (1 - \beta) G(x, y, u'') \subset G(x, y, \beta u' + (1 - \beta) u'');$$

5) limit (6) exists uniformly with respect to u in the domain B ;

6) functions $\varphi(\cdot)$ and $\psi(\cdot)$ are absolutely continuous functions on R_+ and $\varphi(x) \in B', \psi(y) \in B'$ for all $x, y \in R_+$, where $B' + S_\rho(0) \subset B$;

7) the R-solution of the Darboux problem

$$\begin{aligned} u_{xy}^1(x, y) &\in \varepsilon_1 \varepsilon_2 [F(x, y, u^1(x, y))]^1, \\ u^1(x, 0) &= \varphi(x), x \in [0, \infty), \\ u^1(0, y) &= \psi(y), y \in [0, \infty), \varphi(0) = \psi(0), \end{aligned}$$

together with a ρ -neighborhood belong to the domain B for $\varepsilon_1, \varepsilon_2 \in (0, \bar{\varepsilon}]$.

Then for any $\eta \in (0, \rho]$ and $L > 0$ there exists $\varepsilon_0(\eta, L) \in (0, \bar{\varepsilon}]$ such that for all $\varepsilon_1, \varepsilon_2 \in (0, \varepsilon_0]$ and $(x, y) \in [0, L\varepsilon_1^{-1}] \times [0, L\varepsilon_2^{-1}]$ the following inequality holds

$$D(R(x, y), \bar{R}(x, y)) < \eta \quad (7)$$

where $R(\cdot, \cdot), \bar{R}(\cdot, \cdot)$ are the R-solutions of initial and partial averaged Darboux problems.

Proof. By theorem 2, we have unit R-solution of Darboux problem (4) on $[0, L\varepsilon_1^{-1}] \times [0, L\varepsilon_2^{-1}]$ and unit R-solution of

Darboux problem (5) on $[0, L\mathcal{E}_1^{-1}] \times [0, L\mathcal{E}_2^{-1}]$.

Let $\delta_1 = L\mathcal{E}_1^{-1}$, $\delta_2 = L\mathcal{E}_2^{-1}$, $K_n = \left\{ (x_i, y_j) : x_i = ih, y_j = jl, i, j = 0, 1, \dots, n, h = \frac{\delta_1}{n}, l = \frac{\delta_2}{n} \right\}$,

$K_n^{ij} = [x_i, x_{i+1}] \times [y_j, y_{j+1}]$ and $K = [0, L\mathcal{E}_1^{-1}] \times [0, L\mathcal{E}_2^{-1}]$. We denote fuzzy mappings $P^n(\cdot, \cdot)$ and $Q^n(\cdot, \cdot)$ such that

$$[P^n(x, y)]^\alpha = \bigcup_{u \in [P^n(x_i, y_j)]^\alpha} \left\{ v_{ij}(x) + \tau_{ij}(y) - u + \varepsilon_1 \varepsilon_2 \int_{x_i}^x \int_{y_j}^y [F(\zeta, \zeta, u)]^\alpha d\zeta d\zeta \right\},$$

$$[Q^n(x, y)]^\alpha = \bigcup_{z \in [Q^n(x_i, y_j)]^\alpha} \left\{ \bar{v}_{ij}(x) + \bar{\tau}_{ij}(y) - z + \varepsilon_1 \varepsilon_2 \int_{x_i}^x \int_{y_j}^y [G(\zeta, \zeta, z)]^\alpha d\zeta d\zeta \right\},$$

$$P^n(x, 0) = Q^n(x, 0) = \varphi(x), P^n(0, y) = Q^n(0, y) = \psi(y),$$

where $v_{ij}(x) \in AC(x_i, x_{i+1})$, $\tau_{ij}(y) \in AC(y_j, y_{j+1})$, $v_{ij}(x) \in [P^n(x_i, y_j)]^\alpha$, $x \in [x_i, x_{i+1}]$, $\tau_{ij}(y) \in [P^n(x_i, y_j)]^\alpha$, $y \in [y_j, y_{j+1}]$, $v_{ij}(x_i) = \tau_{ij}(y_j) = u$, $\bar{v}_{ij}(x) \in AC(x_i, x_{i+1})$, $\bar{\tau}_{ij}(y) \in AC(y_j, y_{j+1})$, $\bar{v}_{ij}(x) \in [Q^n(x_i, y_j)]^\alpha$, $x \in [x_i, x_{i+1}]$, $\bar{\tau}_{ij}(y) \in [Q^n(x_i, y_j)]^\alpha$, $y \in [y_j, y_{j+1}]$, $\bar{v}_{ij}(x_i) = \bar{\tau}_{ij}(y_j) = z$.

By [52], it follows that the sequences $\{[P^n(\cdot, \cdot)]^\alpha\}_{n=1}^\infty$ and $\{[Q^n(\cdot, \cdot)]^\alpha\}_{n=1}^\infty$ are equicontinuous and fundamental and their limits are α -levels of R-solutions $[R(\cdot, \cdot)]^\alpha$ and $[\bar{R}(\cdot, \cdot)]^\alpha$ of the problems (4) and (5).

Consequently, the sequences $\{P^n(\cdot, \cdot)\}_{n=1}^\infty$ and $\{Q^n(\cdot, \cdot)\}_{n=1}^\infty$ meet by $R(\cdot, \cdot)$ and $\bar{R}(\cdot, \cdot)$.

By [52], for any $\eta_1 > 0$ there exists $0 < \varepsilon_0 \leq \bar{\varepsilon}$ such that

$$h([P^n(x, y)]^\alpha, [Q^n(x, y)]^\alpha) \leq \eta_1 \lambda^{-1} \exp(\lambda L^2), \quad (8)$$

$$h([R(x, y)]^\alpha, [P^n(x, y)]^\alpha) \leq \frac{3\gamma L^2}{n} (1 + \exp(\lambda L^2)), \quad (9)$$

$$h([Q^n(x, y)]^\alpha, [\bar{R}(x, y)]^\alpha) \leq \frac{3\gamma L^2}{n} (1 + \exp(\lambda L^2)) \quad (10)$$

for any $\alpha \in [0, 1]$, $(x, y) \in K$ and $\varepsilon_1, \varepsilon_2 \in (0, \varepsilon_0]$.

Combining (8), (9) and (10), choosing $n \geq \frac{9\gamma L^2 (1 + \exp(\lambda L^2))}{\eta}$ and $\eta_1 < \frac{\eta \lambda}{3 \exp(\lambda L^2)}$ we obtain

$$\begin{aligned} D(R(x, y), \bar{R}(x, y)) &\leq \\ &\leq D(R(x, y), P^n(x, y)) + D(P^n(x, y), Q^n(x, y)) + D(Q^n(x, y), \bar{R}(x, y)) = \\ &= \sup_{\alpha \in [0, 1]} h([R(x, y)]^\alpha, [P^n(x, y)]^\alpha) + \sup_{\alpha \in [0, 1]} h([P^n(x, y)]^\alpha, [Q^n(x, y)]^\alpha) + \\ &\quad + \sup_{\alpha \in [0, 1]} h([Q^n(x, y)]^\alpha, [\bar{R}(x, y)]^\alpha) < \frac{\eta}{3} + \frac{\eta}{3} + \frac{\eta}{3} = \eta. \end{aligned}$$

The theorem is proved.

5. Conclusion.

We conclude with a few remarks.

Remark 1. In this work, we considered the fuzzy differential inclusion, when fuzzy mapping $F(\cdot, u)$ is measurable on $[0, a] \times [0, b]$. If $F(\cdot, u)$ is continuous on $[0, a] \times [0, b]$ then instead of the equation (1) it is possible to consider the following more simple equation

$$\sup_{\alpha \in [0,1]} h([R(t+\sigma, s+\eta)]^\alpha, \bigcup_{u \in [R(t,s)]^\alpha} \{v(t+\sigma) + \tau(s+\eta) - u + \sigma\eta[F(t,s,u)]^\alpha\}) = o(\sigma\eta), \quad (11)$$

and similarly we can prove all results received earlier.

Remark 2. If the condition 4) of Theorem 3 is not true, then the R-solutions can not exist. But there are valid the following conditions:

- 1) for any α -solution $u^\alpha(\cdot, \cdot)$ of inclusion (4) there exists a α -solution $z^\alpha(\cdot, \cdot)$ of inclusion (5) such that $\|u^\alpha(x, y) - z^\alpha(x, y)\| < \eta$ for all $(x, y) \in [0, L\varepsilon_1^{-1}] \times [0, L\varepsilon_2^{-1}]$ and $\alpha \in [0, 1]$;
- 2) for any α -solution $z^\alpha(\cdot, \cdot)$ of inclusion (5) there exists a α -solution $u^\alpha(\cdot, \cdot)$ of inclusion (4) such that $\|u^\alpha(x, y) - z^\alpha(x, y)\| < \eta$ for all $(x, y) \in [0, L\varepsilon_1^{-1}] \times [0, L\varepsilon_2^{-1}]$ and $\alpha \in [0, 1]$.

References

- [1] S. Abbasbandy, T.A. Viranloo, O. Lopez-Pouso, and J. J. Nieto, "Numerical methods for fuzzy differential inclusions," *Computers & Mathematics with Applications*, vol. 48, pp. 1633-1641, 2004. <http://dx.doi.org/10.1016/j.camwa.2004.03.009>.
- [2] N. A. Abdul Rahman, and M.Z. Ahmad, "Applications of the Fuzzy Sumudu Transform for the Solution of First Order Fuzzy Differential Equations," *Entropy*, vol. 17, pp. 4582-4601, 2015.
- [3] A. Arara, M. Benchohra, S. K. Ntouyas, and A. Ouahab, "Fuzzy solutions for hyperbolic partial differential equations," *Int. J. Appl. Math. Sci.*, vol. 2, no. 2, pp. 181-195, 2005.
- [4] R.P. Agarwal, D. O'Regan, and V. Lakshmikantham, "A stacking theorem approach for fuzzy differential equations," *Nonlinear Anal.*, vol. 55, pp. 299-312, 2003, [http://dx.doi.org/10.1016/S0362-546X\(03\)00241-4](http://dx.doi.org/10.1016/S0362-546X(03)00241-4).
- [5] R. P. Agarwal, D. O'Regan, and V. Lakshmikantham, "Maximal solutions and existence theory for fuzzy differential and integral equations," *J. Appl. Anal.*, vol. 11, no. 2, pp. 171-186, 2005, <http://dx.doi.org/10.1515/JAA.2005.171>.
- [6] J.-P. Aubin, "Fuzzy differential inclusions," *Probl. Control Inf. Theory*, vol. 19, no. 1, pp. 55-67, 1990.
- [7] V. A. Baidosov, "Differential inclusions with fuzzy right-hand side," *Sov. Math.*, vol. 40, no. 3, pp. 567-569, 1990.
- [8] V. A. Baidosov, "Fuzzy differential inclusions," *J. of Appl. Math. and Mechan.*, vol. 54, no. 1, pp. 8-13, 1990, [http://dx.doi.org/10.1016/0021-8928\(90\)90080-T](http://dx.doi.org/10.1016/0021-8928(90)90080-T).
- [9] A. M. Bertone, R. M. Jafelice, L. C. de Barros, and R. C. Bassanezi, "On fuzzy solutions for partial differential equations," *Fuzzy Sets Syst.*, vol. 219, pp. 68-80, 2013, <http://dx.doi.org/10.1016/j.fss.2012.12.002>.
- [10] N. N. Bogoliubov, and Yu. A. Mitropolsky, *Asymptotic methods in the theory of non-linear oscillations*. New York, Gordon and Breach, 1961.
- [11] J. J. Buckley, and Th. Feuring, "Introduction to fuzzy partial differential equations," *Fuzzy analysis and related topics* (Prague, 1997). *Fuzzy Sets Syst.*, vol. 105, no. 2, pp. 241-248, 1999, [http://dx.doi.org/10.1016/S0165-0114\(98\)00323-6](http://dx.doi.org/10.1016/S0165-0114(98)00323-6).
- [12] A. Cernea, "On the set of solutions of some nonconvex nonclosed hyperbolic differential inclusions," *Czechoslovak Math. J.*, vol. 52, no. 1, pp. 215-224, 2002, <http://dx.doi.org/10.1023/A:1021787808233>.
- [13] Y. Y. Chen, Y.T. Chang, and B.S. Chen, "Fuzzy Solutions to Partial Differential Equations: Adaptive Approach," *IEEE Transactions on Fuzzy Systems*, vol. 17, no. 1, pp. 116-127, 2009.
- [14] F. S. De Blasi, and J. Myjak, "On the structure of the set of solutions of the Darboux problem of hyperbolic equations," *Proc. Edinburgh Math. Soc. (Series 2)*, vol. 29, no. 1, pp. 7-14, 1986, <http://dx.doi.org/10.1017/S0013091500017351>.
- [15] F. S. De Blasi, and J. Myjak, "On the set of solutions of a differential inclusion," *Bull. Inst. Math. Acad. Sinica*, vol. 14, no. 3, pp. 271-275, 1986.
- [16] F. S. De Blasi, G. Pianigiani, and V. Staicu, "On the solution sets of some nonconvex hyperbolic differential inclusions," *Czechoslovak Math. J.*, vol. 45, no. 120:1, pp. 107-116, 1995.
- [17] D. Dutta Majumder, and K.K. Majumdar, "Complexity analysis, uncertainty management and fuzzy dynamical systems: a cybernetic approach," *Kybernetes*, vol. 33, no. 7, pp. 1143-1184, 2004, <http://dx.doi.org/10.1108/03684920410534489>.
- [18] *Fuzzy partial differential equations and relational equations. Reservoir characterization and modeling*. Edited by Masoud Nikravesh, Lotfi A. Zadeh and Victor Korotkikh in *Studies in Fuzziness and Soft Computing*. vol. 142. Springer-Verlag, Berlin, 2004.
- [19] M. Guo, X. Xue, and R. Li, "Impulsive functional differential inclusions and fuzzy population models," *Fuzzy Sets Syst.*, vol. 138, pp. 601-615, 2003, [http://dx.doi.org/10.1016/S0165-0114\(02\)00522-5](http://dx.doi.org/10.1016/S0165-0114(02)00522-5).
- [20] E. Hüllermeier, "An approach to modelling and simulation of uncertain dynamical system," *Int. J. Uncertain. Fuzziness Knowl.-Based Syst.*, vol. 5, no. 2, pp. 117-137, 1997, <http://dx.doi.org/10.1142/S0218488597000117>.

- [21] E. Hüllermeier, "Numerical methods for Fuzzy Initial Value problems," *Int. J. Uncertain. Fuzziness Knowl.-Based Syst.*, vol. 7, no. 5, pp. 439-461, 1999, <http://dx.doi.org/10.1142/S0218488599000404>.
- [22] R. M. Jafelice, C. G. Almeida, J. F. Meyer, and H. L. Vasconcelos, "Fuzzy parameter in a partial differential equation model for population dispersal of leaf-cutting ants," *Nonlinear Anal. Real World Appl.*, vol. 12, no. 6, pp. 3397-3412, 2011, <http://dx.doi.org/10.1016/j.nonrwa.2011.06.003>.
- [23] M. Kiselevich, "A Bogolyubov type theorem for hyperbolic equations," *Ukrainian Math. J.*, vol. 22, pp. 3, pp. 326-331, 1970, <http://dx.doi.org/10.1007/BF01088955>.
- [24] D.G. Korenevskii, "On the principle of averaging for second-order hyperbolic equations with functionally perturbed argument," *Ukrainian Math. J.*, vol. 23, no. 2, pp. 125-132, 1971, <http://dx.doi.org/10.1007/BF01086602>.
- [25] N. M. Krylov, and N. N. Bogoliubov, *Introduction to nonlinear mechanics*. Princeton, Princeton University Press, 1947.
- [26] V. Lakshmikantham, T. Granna Bhaskar, and J. Vasundhara Devi, *Theory of set differential equations in metric spaces*. Cambridge, UK, Cambridge Scientific Publishers, 2006.
- [27] V. Lakshmikantham, and R. N. Mohapatra, *Theory of fuzzy differential equations and inclusions*. London, UK, Taylor & Francis, 2003.
- [28] A. Lomtadze, and J. Šremr, "Caratheodory solutions to a hyperbolic differential inequality with a non-positive coefficient and delayed arguments," *Bound. Value Probl.*, vol. 2014:52, pp. 1-13, 2014, <http://dx.doi.org/10.1186/1687-2770-2014-52>.
- [29] V. L. Lažar, "Ulam-Hyers stability for partial differential inclusions," *Electron. J. Qual. Theory Differ. Equ.*, vol. 21, pp. 1-19, 2012, <http://dx.doi.org/10.14232/ejqtde.2012.1.21>.
- [30] H. V. Long, N. T. K. Son, N. T. M. Ha, and L. H. Son, "The existence and uniqueness of fuzzy solutions for hyperbolic partial differential equations," *Fuzzy Optimization and Decision Making*, vol. 13, no. 4, pp. 435-462, 2014, <http://dx.doi.org/10.1007/s10700-014-9186-0>.
- [31] C. V. Negoita, and D.A. Ralescu, *Application of fuzzy sets to systems analysis*. New York, USA, Wiley, 1975.
- [32] Norazrizal Aswad Abdul Rahman, and Muhammad Zaini Ahmad, "Fuzzy Sumudu transform for solving fuzzy partial differential equations," *J. Nonlinear Sci. Appl.*, vol. 9, pp. 3226-3239, 2016.
- [33] Nayyar Mehmood, and Akbar Azam, "Existence Results for Fuzzy Partial Differential Inclusions," *Journal of Function Spaces*, Article ID 6759294, 8 pages, 2016, <http://dx.doi.org/10.1155/2016/6759294>.
- [34] A. I. Panasyuk, and V.I. Panasyuk, "An equation generated by a differential inclusion," *Mathematical notes of the Academy of Sciences of the USSR*, vol. 27, no. 3, pp. 213-218, 1980, <http://dx.doi.org/10.1007/BF01140170>.
- [35] U. M. Pirzada, and D.C. Vakaskar, "Solution of fuzzy heat equations using Adomian Decomposition method," *Int. J. Adv. Appl. Math. and Mech.*, vol. 3, no. 1, pp. 87-91, 2015.
- [36] A. V. Plotnikov, "A procedure of complete averaging for fuzzy differential inclusions on a finite segment," *Ukrainian Math. J.*, vol. 67, no. 3, pp. 421-430, 2015, <http://dx.doi.org/10.1007/s11253-015-1090-4>.
- [37] A. V. Plotnikov, and T. A. Komleva, "The partial averaging of fuzzy differential inclusions on finite interval," *International Journal of Differential Equations*, Article ID 307941, 5 pages, 2014, <http://dx.doi.org/10.1155/2014/307941>.
- [38] A. V. Plotnikov, and T. A. Komleva, "The Averaging Of Fuzzy Linear Differential Inclusions On Finite Interval," *Dynamics of Continuous, Discrete and Impulsive Systems, Series B: Applications & Algorithms*, vol. 23, no. 1, pp. 1-9, 2016.
- [39] A. V. Plotnikov, T. A. Komleva, and L.I. Plotnikova, "The partial averaging of differential inclusions with fuzzy right-hand side," *J. Adv. Res. Dyn. Control Syst.*, vol. 2, no. 2, pp. 26-34, 2010.
- [40] V. A. Plotnikov, A. V. Plotnikov, and A. N. Vityuk, *Differential equations with a multivalued right-hand side. Asymptotic methods*. Odessa, Ukraine, AstroPrint, 1999.
- [41] M. L. Puri, and D. A. Ralescu, "Fuzzy random variables," *J. Math. Anal. Appl.*, vol. 114, pp. 409-422, 1986, [http://dx.doi.org/10.1016/0022-247X\(86\)90093-4](http://dx.doi.org/10.1016/0022-247X(86)90093-4).
- [42] J. A. Sanders, and F. Verhulst, *Averaging methods in nonlinear dynamical systems*. Appl. Math. Sci., vol. 59, New York, Springer-Verlag, 1985.
- [43] J. Šremr, "Absolutely continuous functions of two variables in the sense of Caratheodory," *Electron. J. Diff. Equ.*, vol. 2010, no. 154, pp. 1-11, 2010.
- [44] V. Staicu, "On a non-convex hyperbolic differential inclusion," *Proc. Edinburgh Math. Soc. (Series 2)*, vol. 35, no. 3, pp. 375-382, 1992, <http://dx.doi.org/10.1017/S0013091500005666>.
- [45] G. Teodoru, "Approximation of the solution of Darboux problem for the equation $\partial^2 z / \partial x \partial y \in F(x, y, z)$," *Itinerant seminar on functional equations, approximation and convexity (Cluj-Napoca, 1985)*. Preprint 85-6. Univ. "Babes-Bolyai", Cluj-Napoca, pp. 215-220, 1985.
- [46] G. Teodoru, "On the Darboux problem for the equation $\partial^2 z / \partial x \partial y \in F(x, y, z)$," *An. Stiint. Univ. Al. I. Cuza Iasi Sect. I a Mat.*, vol. 32, no. 3, pp. 41-49, 1986.
- [47] G. Teodoru, "Continuous selections for multifunctions satisfying the Caratheodory type conditions. The Darboux problem associated to a multivalued equation," *Itinerant Seminar on Functional Equations, Approximation and Convexity (Cluj-Napoca, 1987)*. Preprint 87-6. Univ. "Babes-Bolyai", Cluj-Napoca, pp. 281-286, 1987.
- [48] G. Teodoru, "A characterization of the solutions of the Darboux problem for the equation $\partial^2 z / \partial x \partial y \in F(x, y, z)$," *An. Stiint. Univ. Al. I. Cuza Iasi Mat.*, vol. 33, no. 1, pp. 33-38, 1987.
- [49] A. N. Vityuk, "Properties of solutions of hyperbolic differential equations with many-valued right-hand sides," *Mat. Fiz. Nelin. Mekh.*, vol. 15, pp. 59-62, 1991.
- [50] A. N. Vityuk, "Equation of the integral funnel of a partial differential inclusion," *Dokl. Ukr. Akad. Nauk, Ser. A.*, vol. 9, pp. 19-20, 1992.

- [51] A. N. Vityuk, "On solutions of hyperbolic differential inclusions with nonconvex right-hand side," *Ukrainian Math. J.*, vol. 47, no. 4, pp. 617-621, 1995, <http://dx.doi.org/10.1007/BF01056048>.
- [52] A. N. Vityuk, "On an R-solution generated by a differential inclusion of hyperbolic type," *Differential Equations*, vol. 30, no. 10, pp. 1578-1586, 1994.
- [53] A. N. Vityuk, "Continuous dependence of the R-solution generated by a differential hyperbolic inclusion on parameters," *Ukrainian Math. J.*, vol. 47, no. 10, pp. 1625-1631, 1995, <http://dx.doi.org/10.1007/BF01060163>.